CDM QIP indicators for revised flowsheets to be released in 2024.

Changes from 2016 document highlighted in yellow

DIABETES (DM)		
CDM indicators	CDM-QIP Indicator description (Indication of optimal clinical practice)	What the payment system looks for in a 12-month period for each patient (Based on indicator data exported from flowsheets or entered into EHR viewer)
Type of diabetes	Record whether patient has "type 1" or "type 2" or "other"	Type of diabetes needs to be noted in patient record and CDMQIP flowsheet at least once for payment
Blood pressure	Measurement of BP and recorded at least twice in 12 months	Presence of systolic and diastolic BP values on 2 separate dates
Monitoring of weight	Measurement of body weight at least once in 12 months (Height and BMI no longer required)	Presence of one weight value
Smoking status and smoking cessation advice	Smoking status reviewed and recorded at least twice in 12 months; cessation advice provided to smokers at least twice in 12 months	Presence of smoking status on 2 separate dates and presence of smoking cessation advice for patients who are smokers
Glycemic control	A1c performed and value recorded at least twice in 12 months	Presence of A1c value and test dates on 2 separate dates
Lipid screening	Lipid profile at least annually/as needed (testing influenced by age, duration of diabetes, CVD risk factors, treatment). LDL value recorded at least once in 12 months for patients aged 40-80 years	If patient age is 40-80 years, presence of 1 LDL cholesterol value and lipid test date
Nephropathy screening and monitoring	Urine ACR and eGFR recorded at least once in 12 months, and nephropathy status recorded (ACR not required if end-stage renal disease eGFR <15 mL/min)	Presence of 5 components: Urine ACR value and date of test, eGFR value and date of test, and nephropathy status. If eGFR value <15 mL/min the presence of 3 components is required: EGFR value and date and nephropathy status
Retinopathy screening and monitoring	Diabetic retinopathy status recorded at least once in 12 months	Presence of retinopathy status
Foot exam and peripheral neuropathy screening and monitoring	Diabetic foot exam completed at least once in 12 months, and peripheral neuropathy status recorded	Presence of diabetic foot exam and date and neuropathy status
Depression/psychosocial stress screening **	Depression/psychosocial stress evaluation performed at least once in 12 months	Presence of depression screening date and response of "Yes" or "No" to "experiencing low mood/anhedonia/anxiety"

** Screening for depression/stress has been modified; HCP now responds to the question "Patient experiencing low mood/anhedonia/anxiety?" with "Yes" or "No". If response left blank or "Not asked" checked, the indicator will not be considered for QIP payment.