

HEART FAILURE (HF)		
CDM indicators	CDM-QIP Indicator description (Indication of optimal clinical practice)	What the payment system looks for in a 12-month period for each patient (Based on indicator data exported from flowsheets or entered into EHR viewer)
Type of heart failure	Record whether patient has HFrEF, HFpEF or HFmEF	Type of HF needs to be noted in patient record and CDMQIP flowsheet <i>at least once for payment</i>
Assessment of HF functional status	NYHA functional class assessed and recorded at least twice in 12 months	Presence of NYHA functional class value on 2 separate dates
Blood pressure	Measurement of BP and recorded at least twice in 12 months	Presence of systolic and diastolic BP values on 2 separate dates
Volume assessment	Volume /fluid status assessed and recorded at least twice in 12 months	Presence of volume status response (euvolemic, hypovolemic, fluid overload) on 2 separate dates
Monitoring of weight	Measurement of body weight at least twice in 12 months	Presence of weight value on 2 separate dates
Smoking status and smoking cessation advice	Smoking status reviewed and recorded at least twice in 12 months; cessation advice provided to smokers at least twice in 12 months	Presence of smoking status on 2 separate dates and presence of smoking cessation advice for patients who are smokers
Diabetes screening for patients who do not have diabetes	Fasting blood glucose or A1c tests performed at least once in 12 months	If the patient exists in the payment system database as already having diabetes, this indicator is ignored. If the patient does not have diabetes “flag” in the payment system, 1 fasting glucose value and test date or 1 A1c value and test date must be present
Obstructive sleep apnea (OSA) screening	Screen for OSA at least once in 12 months	Presence of 1 of the OSA screening responses
ACE inhibitor or ARB or ARNI therapy/use	ACE inhibitor, ARB or ARNI therapy/use assessed and recorded at least once in 12 months	Presence of 1 of the ACE inhibitor, ARB or ARNI therapy responses
Beta-blocker therapy/use	Beta-blocker therapy/use assessed and recorded at least once in 12 months	Presence of 1 of the beta-blocker therapy responses
MRA therapy/use	MRA therapy/use assessed and recorded at least once in 12 months	Presence of 1 of the MRA therapy responses
Depression/psychosocial stress screening **	Depression/psychosocial stress evaluation performed at least once in 12 months	Presence of depression screening date and response of “Yes” or “No” to “experiencing low mood/anhedonia/anxiety”

** Screening for depression/stress has been modified; HCP now responds to the question “Patient experiencing low mood/anhedonia/anxiety?” with “Yes” or “No”. If response left blank or “Not asked” checked, the indicator will not be considered for QIP payment.