Immunization of immunocompromised adults and children with live attenuated varicella vaccine <u>requires approval</u> from the medical specialist, primary care physician or nurse practitioner most familiar with the client's current medical status.		
Patient name: Gender HSN		
DOB:	_ Gender HSN	l
Check the appropriate condition below:		
HIV infection ¹ – only individuals who are not severely immunosuppressed as indicated according to criteria based on age for immunization:		 Isolated immune deficiency² Humoral (Ig) deficiency diseases Neutrophil deficiency diseases
Age 1 through 5 years old	Age ≥ 6 years old	Complement deficiency diseases
CD4+ T-lymphocyte counts (x10 ⁶ /L) for ≥ 6 months: ≥ 500	CD4+ T-lymphocyte counts (x10 ⁶ /L) for ≥ 6 months: ≥ 200	 □ Transplant recipient □ Kidney □ Liver □ HSCT □ Other (specify) □ Long-term immunosuppressive therapy³ □ Other condition (specify):
Total lymphocytes: ≥ 15%	Total lymphocytes: ≥15%	Comments:
1) If applicable: VZV IgG test result: Date of test:		
 I verify on that this patient has no medical contraindications to receiving varicella vaccine. 		
3) I understand that this patient may require up to 2 doses given at least 3 months apart <u>and</u> verify that this client's condition is sufficiently stable to receive up to 2 doses within an 8-month period from when this referral has been received at Public Health.		
		1D DNP Clinic:
Phone #:	Fax #:	Email:
Dose 1 date given:	Lot #:	
Dose 2 date given:		
Public Health Nurse's name (print):		
Phone #:	Fax:	Email:

¹ Varicella and MMR may be administered on same day. MMRV vaccine contraindicated.

² Min. 4-week intervals are required between all Varicella and MMR vaccine doses. MMRV vaccine contraindicated.

³ Live vaccines are contraindicated during immunosuppressive therapy. However, the patient's clinical status, net state of immune suppression (including immunosuppressive medications and presence of opportunistic infections), and an indepth immunologic evaluation are important considerations when deciding about risk versus benefit for live virus immunization.

